

On The Relevance of Fair Sampling Assumption in The Recent Bell Photonic Experiments

A. Shafiee^{*(1)} and M. Golshani^(2,3)

1) Department of Chemistry, Sharif University of Technology,
P.O.Box 11365-9516, Tehran, Iran

2) Department of Physics, Sharif University of Technology,
P.O.Box 11365-9161, Tehran, Iran.

3) Institutes for Studies in Theoretical Physics & Mathematics,
P.O.Box 19395-5531, Tehran, Iran.

February 1, 2008

Abstract

In the experimental verification of Bell's inequalities in real photonic experiments, it is generally believed that the so-called fair sampling assumption (which means that a small fraction of results provide a fair statistical sample) has an unavoidable role. Here, we want to show that the interpretation of these experiments could be feasible, if some different alternative assumptions other than the fair sampling were used. For this purpose, we derive an efficient Bell-type inequality which is a CHSH-type inequality in real experiments. Quantum mechanics violates our proposed inequality, independent of the detection-efficiency problems.

1 Introduction

In his celebrated 1964 paper, John Bell considered a system consisting of two spin- $\frac{1}{2}$ particles in a singlet state [1]. He showed that the correlation

^{*}E-mail: shafiee@sharif.edu

between the results of two experiments done on such spatially separated particles cannot be reproduced by a local hidden-variable theory. Experiments done since 1972 indicate that the spin correlations of two particles in a singlet state violate Bell's inequality, as quantum mechanics requires. These experiments have been usually done for photons (see ref. [2] and the references therein), and recently they were performed for massive particles [3]. But, there has been two general loopholes in the standard interpretation of these experiments which are not yet taken care of simultaneously in a single experiment. They make the present interpretations inconclusive. These are known as detection loophole and locality loophole.

The detection loophole [4] refers to the fact that in Bell-type experiments, due to the low efficiency of detectors and collimators, a large number of photons may be undetected, and the resulting correlation is obtained on the basis of detected photons. Consequently, it is always possible to construct a local hidden variable model which can reproduce the experimental results [4, 5, 6]. So, the interpretation of these experiments is only feasible if one makes the fair sampling (FS) assumption [7]. P. Grangier describes the detection-efficiency loophole as "Achill's heel of experimental tests of Bell's inequalities" [8]. In the experiments done with the massive particles, this problem is solved, because the detection of these particles could be done efficiently. Yet, the second loophole, i.e., the locality loophole (according to which there exist the possibility of (sub)luminal communication between two spatially separated particles) is still there. (see, e.g., ref. [9].)

In this paper, we consider the problem of fair sampling in the experiments done with photons. (A complete review of these experiments can be found in ref. [2].) There have been considerable discussions in the literature on this subject. But, the main issue in all of them is either to obtain a new limit for the detector inefficiency in CH [10] and/or CHSH [7] inequalities (see, e.g., ref. [11]) or to include the detector inefficiency directly into the original Bell inequality (see, e.g., ref. [12]). Recently, the authors have also proposed a new CH-type inequality which can be violated by some quantum mechanical predictions independent of the efficiency factors [13, 14].

What we want to argue here is that, contrary to what is assumed so far, detection loophole could be neglected in the interpretations of the recent photonic experiments, since there are other independent assumptions which are reasonable by themselves and could be used. To show this, we use, in part 2 of this paper, an *efficient* type of CHSH inequality which has been tested in recent photonic experiments and is shown to have been violated. We argue that for observing the violation of this inequality in real experiments, there are at least three possible solutions (other than the FS assumption) which have different physical basis and can be used independently for deriving

the inequality. In section 3, we show that quantum mechanical predictions violate this inequality for real experiments, *independent* of the efficiency of detectors and collimators. Thus, we find *another* way for the justification of the entanglement criteria in the microphysical Bell states.

2 An Alternative Bell-type Inequality

FS assumption means that unrecorded data *do not* have a weighty role in calculating the polarization correlations of the two entangled photons. This is the most common view about the FS assumption. That is what P. Pearle described as Data Rejection Hypothesis in his 1970 paper [4]: “Suppose that each particle has three responses to a spin-measuring apparatus instead of two....Then, instead of four possible experimental outcomes of the measurement of the spins of two particles, there are nine possible outcomes. In one of these outcomes, neither particle is detected, and so the experimenter is unaware that a decay has taken place. In four of these outcomes one of the particles is not detected. If the experimenter rejects these data (in the belief that the apparatus is not functioning properly and that if it had been functioning properly, the data recorded would have been representative of the accepted data), he is left with the usual four possible outcomes.” Similarly, P. Grangier describes the meaning of fair sampling assumption as [8]: “The detection-efficiency loophole argues that, in most experiments, only a very small fraction of the particles generated are actually detected....So, to extract a meaningful conclusion from the observed data, it was necessary to assume that a small fraction of data provides a fair statistical sample”. Considering FS assumption, however, *there is no reason why the data recorded are representative of the accepted data, what is the nature of recording probabilities and how one can interpret the efficiencies*. These questions are *irrelevant* when one refers to the FS assumption. In contrast, if we negate such a hypothesis, i.e. if we believe that the rejected data may have a significant role in calculating the correlations, it would be a crucial task to elucidate the above points. This is our main concern in the following. (See the appendix for a more concrete discussion about the FS assumption.)

Let us consider an actual double-channel Bell experiment where for each emitted photon a binary event (i.e., passage or non-passage corresponding to two polarization eigenvalues) occurs at each analyzer. We assume a stochastic local hidden-variable (SLHV) theory, in which λ represents a collection of hidden variables, belonging to a space Λ . To have a complete physical description of the whole system, the hidden variables λ are assumed to include the underlying variables of both the particles and devices. At this level,

$p_r^{(1)}(\hat{a}, \lambda)$ is the probability that the result r is detected for the polarization of the first photon along $\hat{a}$, where $r = \pm 1$ corresponds to two eigenvalues of photon's polarization and the angle $\hat{a}$ is an angle from the x -axis to the transmission axis of the first photon's polarization filter. The detection probability $p_r^{(1)}(\hat{a}, \lambda)$ can be defined as the following:

$$p_r^{(1)}(\hat{a}, \lambda) = p_{r,id}^{(1)}(\hat{a}, \lambda) \eta_{1r}(\hat{a}, \lambda) \quad (1)$$

Here, $p_{r,id}^{(1)}(\hat{a}, \lambda)$ is the probability that if the first photon encounters a polarizer at angle $\hat{a}$, it will then be detected in channel r in an *ideal* experiment. In an analogous real experiment, we assume that $\eta_{1r}(\hat{a}, \lambda)$ denotes the overall efficiency of detecting the first photon with polarization along $\hat{a}$ in channel r . It contains, e.g., the probability that the first photon reaches its detector and then will be detected with a definite chance. One can define $p_q^{(2)}(\hat{b}, \lambda)$ in a similar fashion for the second photon with $q = \pm 1$.

The probability of non-detection of photons 1 and 2, along $\hat{a}$ and $\hat{b}$ respectively, are represented by $p_0^{(1)}(\hat{a}, \lambda)$ and $p_0^{(2)}(\hat{b}, \lambda)$, where the index zero denotes non-detection. According to relation (1), $p_0^{(1)}(\hat{a}, \lambda)$ can be defined as $p_0^{(1)}(\hat{a}, \lambda) = 1 - \alpha(\hat{a}, \lambda)$, where $\alpha(\hat{a}, \lambda) = \sum_{r=\pm 1} p_r^{(1)}(\hat{a}, \lambda)$ is a representative function of the overall hidden efficiency $\eta_{1r}(\hat{a}, \lambda)$. If one assumes that $\eta_{1r}(\hat{a}, \lambda)$ is independent of the measured value of the polarization r (which means that all the efficiencies are the same at two channels $+$ and $-$), then $\alpha(\hat{a}, \lambda) = \eta_1(\hat{a}, \lambda)$ can be interpreted as an overall measure of the efficiency at the hidden-variable level. I.e.,

$$p_0^{(1)}(\hat{a}, \lambda) = 1 - \eta_1(\hat{a}, \lambda) \quad (2)$$

A similar relation can be considered for $p_0^{(2)}(\hat{b}, \lambda)$.

The joint probability for detection of the two photons with the outcomes r and q corresponding to the polarizations along $\hat{a}$ and $\hat{b}$, respectively, is assumed to be:

$$p_{rq}^{(12)}(\hat{a}, \hat{b}, \lambda) = p_r^{(1)}(\hat{a}, \lambda) p_q^{(2)}(\hat{b}, \lambda) \quad (3)$$

This is known as Bell's locality condition [10]. Relations similar to (3) hold for the joint probabilities concerning non-detections.

In a SLHV theory, the average value of the outcomes of polarizations of two photons along $\hat{a}$ and $\hat{b}$ is given by

$$\varepsilon^{(12)}(\hat{a}, \hat{b}, \lambda) = \sum_{r,q=\pm 1} r q p_{rq}^{(12)}(\hat{a}, \hat{b}, \lambda)$$

$$= \varepsilon^{(1)}(\hat{a}, \lambda) \varepsilon^{(2)}(\hat{b}, \lambda) \quad (4)$$

where $\varepsilon^{(1)}(\hat{a}, \lambda)$ and $\varepsilon^{(2)}(\hat{b}, \lambda)$ are the average values of the outcomes of polarizations for photons 1 and 2 along $\hat{a}$ and $\hat{b}$, respectively. Assuming that the above probabilities are normalized to one, we have:

$$\sum_{j=\pm 1,0} p_j^{(1)}(\hat{a}, \lambda) = \sum_{j=\pm 1,0} p_j^{(2)}(\hat{b}, \lambda) = 1 \quad (5)$$

Now, it is obvious that

$$0 \leq p_r^{(1)}(\hat{a}, \lambda) \leq \alpha(\hat{a}, \lambda) \quad (6)$$

and

$$0 \leq p_q^{(2)}(\hat{b}, \lambda) \leq \beta(\hat{b}, \lambda) \quad (7)$$

where $\alpha(\hat{a}, \lambda) = \sum_{r=\pm 1} p_r^{(1)}(\hat{a}, \lambda)$ and $\beta(\hat{b}, \lambda) = \sum_{q=\pm 1} p_q^{(2)}(\hat{b}, \lambda)$. The constraints (6) and (7) are actual constraints for the detection of single particles at the hidden-variable level. In the ideal limit, where $\alpha(\hat{a}, \lambda) \rightarrow 1$ and $\beta(\hat{b}, \lambda) \rightarrow 1$, the probability of detection lies within the interval $[0, 1]$. Using the aforementioned constraints, one gets:

$$| \varepsilon^{(1)}(\hat{a}, \lambda) | \leq \alpha(\hat{a}, \lambda) \quad (8)$$

and

$$| \varepsilon^{(2)}(\hat{b}, \lambda) | \leq \beta(\hat{b}, \lambda) \quad (9)$$

In the following, we introduce three independent solutions which include some plausible assumptions about the nature of *non-detection* probabilities at the level of hidden variables as well as the relation of the empirical correlations with the predictions of a SLHV theory. These assumptions provide alternative ways for deriving an *efficient* type of CHSH inequality. Then, one can argue for the soundness of the recent photonic experiments. Yet, there are some important points which should be noted here. Our solutions I and II below involve assumptions about the probabilities of non-detection. The non-detection probabilities are unobservable and it has been usually recommended to avoid them. Thus, the earlier works in this area involved constraints about the probabilities of detection, rather than non-detection [7, 10]. Nevertheless, it is reasonable to think that a more plausible approach with *weaker* assumptions is achieved when one takes into account the non-detection events. This is the main point of the present work in which the

nature of the auxiliary assumptions are completely different with the so-called fair sampling or no-enhancement assumptions in CHSH or CH inequalities.

As indicated before, what we are proposing here is that the non-detection probabilities *do* have an important role in calculating the photonic correlations. But, we shall argue that there are situations in which one can define an *effective* correlation function only based on *detected* events and derive an inequality which only contains the so-called effective correlations. This is our purposed inequality. Here, we survey these situations in the context of the following solutions.

Solution I. This is based on the assumption that *at the level of hidden variables, the probability of non-detection of each individual photon is independent of the direction of its polarization filter*, i.e.,

$$p_0^{(1)}(\hat{a}, \lambda) = p_0^{(1)}(\hat{a}', \lambda); \quad (10-a)$$

$$p_0^{(2)}(\hat{b}, \lambda) = p_0^{(2)}(\hat{b}', \lambda) \quad (10-b)$$

According to relation (1), this means also that for each individual photon, the hidden probabilities for reaching a detector and detecting by it should be independent of the earlier preparation made by choosing a definite polarization angle.

Now, let us consider the set of polarization directions $\hat{a}, \hat{a}'$ for the first photon and $\hat{b}, \hat{b}'$ for the second one. Furthermore, we define the function u as

$$u := x(y - y') + x'(y + y') \quad (11)$$

where $x := \varepsilon^{(1)}(\hat{a}, \lambda)$, $x' := \varepsilon^{(1)}(\hat{a}', \lambda)$, $y := \varepsilon^{(2)}(\hat{b}, \lambda)$ and $y' := \varepsilon^{(2)}(\hat{b}', \lambda)$. We have also $|x| \leq \alpha$, $|x'| \leq \alpha'$, $|y| \leq \beta$ and $|y'| \leq \beta'$, in which for example $\alpha := \alpha(\hat{a}, \lambda) = 1 - p_0^{(1)}(\hat{a}, \lambda)$, $\beta := \beta(\hat{b}, \lambda) = 1 - p_0^{(2)}(\hat{b}, \lambda)$ and similar definitions hold for α' and β' . Considering the relations (10-a) and (10-b), we have $\alpha = \alpha' = \alpha(\lambda)$ and $\beta = \beta' = \beta(\lambda)$. So, the limits of $|x|$ and $|x'|$ as well as $|y|$ and $|y'|$ are the same. Since u is a linear function of the variables x, x', y and y' , its upper and lower bounds are determined by the limits of these variables. The bounds are tabulated in the Table 1. This table shows that u is confined by the limits $2\alpha\beta$ and $-2\alpha\beta$.

Rows	x	x'	y	y'	u
1	$-\alpha$	$-\alpha'$	$-\beta$	$-\beta'$	$\alpha(\beta - \beta') + \alpha'(\beta + \beta') = 2\alpha\beta$
2	α	$-\alpha'$	$-\beta$	$-\beta'$	$-\alpha(\beta - \beta') + \alpha'(\beta + \beta') = 2\alpha\beta$
3	$-\alpha$	α'	$-\beta$	$-\beta'$	$\alpha(\beta - \beta') - \alpha'(\beta + \beta') = -2\alpha\beta$
4	$-\alpha$	$-\alpha'$	β	$-\beta'$	$-\alpha(\beta + \beta') - \alpha'(\beta - \beta') = -2\alpha\beta$
5	$-\alpha$	$-\alpha'$	$-\beta$	β'	$\alpha(\beta + \beta') + \alpha'(\beta - \beta') = 2\alpha\beta$
6	α	α'	$-\beta$	$-\beta'$	$-\alpha(\beta - \beta') - \alpha'(\beta + \beta') = -2\alpha\beta$
7	α	$-\alpha'$	β	$-\beta'$	$\alpha(\beta + \beta') - \alpha'(\beta - \beta') = 2\alpha\beta$
8	α	$-\alpha'$	$-\beta$	β'	$-\alpha(\beta + \beta') + \alpha'(\beta - \beta') = -2\alpha\beta$
9	$-\alpha$	α'	β	$-\beta'$	$-\alpha(\beta + \beta') + \alpha'(\beta - \beta') = -2\alpha\beta$
10	$-\alpha$	α'	$-\beta$	β'	$\alpha(\beta + \beta') - \alpha'(\beta - \beta') = 2\alpha\beta$
11	$-\alpha$	$-\alpha'$	β	β'	$-\alpha(\beta - \beta') - \alpha'(\beta + \beta') = -2\alpha\beta$
12	α	α'	β	$-\beta'$	$\alpha(\beta + \beta') + \alpha'(\beta - \beta') = 2\alpha\beta$
13	α	α'	$-\beta$	β'	$-\alpha(\beta + \beta') - \alpha'(\beta - \beta') = -2\alpha\beta$
14	α	$-\alpha'$	β	β'	$\alpha(\beta - \beta') - \alpha'(\beta + \beta') = -2\alpha\beta$
15	$-\alpha$	α'	β	β'	$-\alpha(\beta - \beta') + \alpha'(\beta + \beta') = 2\alpha\beta$
16	α	α'	β	β'	$\alpha(\beta - \beta') + \alpha'(\beta + \beta') = 2\alpha\beta$

Table 1: The limits of u .

Thus, under these conditions, we have:

$$|u| \leq 2\alpha\beta \quad (12)$$

In the ideal limit we have $|u| \leq 2$.

Now, we assume that the empirical correlation functions have a definite relation with the averages of the outcomes of polarizations of the two photons along certain directions in a SLHV theory. For example, for the two polarization directions $\hat{a}$ and $\hat{b}$, we define:

$$E^{(12)}(\hat{a}, \hat{b}) = \int_{\Lambda} \varepsilon^{(1)}(\hat{a}, \lambda) \varepsilon^{(2)}(\hat{b}, \lambda) \rho(\lambda) d\lambda \quad (13)$$

where, $E^{(12)}(\hat{a}, \hat{b})$ is the correlation function of the polarization measurements of the two photons along $\hat{a}$ and $\hat{b}$, and $\rho(\lambda)$ is the normalized probability density of λ over Λ . Using the definitions of α and β , we have:

$$\sum_{r,q=\pm 1} P_{rq}^{(12)} = \int_{\Lambda} \alpha(\lambda) \beta(\lambda) \rho(\lambda) d\lambda \quad (14)$$

where $P_{rq}^{(12)}$ is the probability of the simultaneous detection of the outcome r for the first photon and q for the second photon, with polarizations along two arbitrary directions, in a real experiment. The relation (14) is independent

of the polarization directions. But, this does not mean that the total number of photons recorded by each detector is independent of the directions of the polarization filters, because the number of undetected photons has a weighty role in the definition of the detection probabilities.

Using the relations (11), (13) and (14), the inequality (12) takes the following form:

$$|U| \leq M \quad (15)$$

where

$$U = E^{(12)}(\hat{a}, \hat{b}) - E^{(12)}(\hat{a}, \hat{b}') + E^{(12)}(\hat{a}', \hat{b}) + E^{(12)}(\hat{a}', \hat{b}') \quad (16)$$

and $M = 2 \sum_{r,q=\pm 1} P_{rq}^{(12)}$.

In an ideal case, we have $M = 2$ where by ideal we mean an experiment in which the probabilities of non-detection are zero. In view of the fact that in general $M \leq 2$, one can infer from (15) that

$$|U| \leq 2 \quad (17)$$

The inequality (17) is known as CHSH inequality in the literature.

Now, we define the effective correlation functions measured in the photonic experiments as

$$E_{eff}^{(12)}(\hat{a}, \hat{b}) := \frac{E^{(12)}(\hat{a}, \hat{b})}{\sum_{r,q=\pm 1} P_{rq}^{(12)}(\hat{a}, \hat{b})} = \frac{\sum_{r,q=\pm 1} rq N_{rq}^{(12)}(\hat{a}, \hat{b})}{\sum_{r,q=\pm 1} N_{rq}^{(12)}(\hat{a}, \hat{b})} \quad (18)$$

where $N_{rq}^{(12)}(\hat{a}, \hat{b})$ is the number of photons that are detected with the outcomes r and q along $\hat{a}$ and $\hat{b}$, respectively. Assuming that $P_{00}^{(12)}(\hat{a}, \hat{b}) = P_0^{(1)}(\hat{a})P_0^{(2)}(\hat{b})$, we have $\sum_{r,q=\pm 1} P_{rq}^{(12)}(\hat{a}, \hat{b}) = (1 - P_0^{(1)}(\hat{a}))(1 - P_0^{(2)}(\hat{b}))$, where $P_0^{(1)}(\hat{a})$ ($P_0^{(2)}(\hat{b})$) is the probability of non-detection of photon 1 (2) with a polarization along $\hat{a}$ ($\hat{b}$) and $P_{00}^{(12)}(\hat{a}, \hat{b})$ is the joint probability of non-detection for both photons. Using the definition (18), the inequality (15) is reduced to

$$|U_{eff}| \leq 2 \quad (19)$$

where

$$U_{eff} = E_{eff}^{(12)}(\hat{a}, \hat{b}) - E_{eff}^{(12)}(\hat{a}, \hat{b}') + E_{eff}^{(12)}(\hat{a}', \hat{b}) + E_{eff}^{(12)}(\hat{a}', \hat{b}') \quad (20)$$

The inequality (19) is our proposed Bell-type inequality for a real experiment. This is the inequality which has been tested in recent photonic experiments and is shown to have been violated.

Solution II. The first solution was based on assumptions that are used at the hidden-variable level. In the second solution, however, both the experimental and hidden-variable levels are under consideration. To derive (19), we assume that *non-detection probabilities for each individual photon are the same at both levels*, i.e.,

$$P_0^{(1)}(\hat{a}) = p_0^{(1)}(\hat{a}, \lambda); \quad P_0^{(1)}(\hat{a}') = p_0^{(1)}(\hat{a}', \lambda) \quad (21-a)$$

$$P_0^{(2)}(\hat{b}) = p_0^{(2)}(\hat{b}, \lambda); \quad P_0^{(2)}(\hat{b}') = p_0^{(2)}(\hat{b}', \lambda) \quad (21-b)$$

Here, one can argue that non-detection probabilities are hidden, as is the case at the hidden-variable level. Because, there is no way for their detection. The necessary condition for the acceptance of above relations is the assumption that the non-detection probability for each individual photon, at the hidden-variable level, is independent of λ . Or, equivalently, this means that the hidden efficiencies for reaching a detector and detection by it are the same as the experimental ones (see relation (1)).

Subsequently, One can define an effective average value at the level of hidden variables, as

$$\begin{aligned} \varepsilon_{eff}^{(12)}(\hat{a}, \hat{b}, \lambda) &= \sum_{r,q=\pm 1} r q \left(\frac{p_r^{(1)}(\hat{a}, \lambda)}{1 - p_0^{(1)}(\hat{a}, \lambda)} \right) \left(\frac{p_q^{(2)}(\hat{b}, \lambda)}{1 - p_0^{(2)}(\hat{b}, \lambda)} \right) \\ &= \varepsilon_{eff}^{(1)}(\hat{a}, \lambda) \varepsilon_{eff}^{(2)}(\hat{b}, \lambda) \end{aligned} \quad (22)$$

where $\varepsilon_{eff}^{(1)}(\hat{a}, \lambda) = \sum_{r=\pm 1} r \left(\frac{p_r^{(1)}(\hat{a}, \lambda)}{1 - p_0^{(1)}(\hat{a}, \lambda)} \right)$ and $\varepsilon_{eff}^{(2)}(\hat{b}, \lambda) = \sum_{q=\pm 1} q \left(\frac{p_q^{(2)}(\hat{b}, \lambda)}{1 - p_0^{(2)}(\hat{b}, \lambda)} \right)$. Using (6) and (7), we get:

$$\left| \varepsilon_{eff}^{(1)}(\hat{a}, \lambda) \right| \leq 1 \quad (23)$$

$$\left| \varepsilon_{eff}^{(2)}(\hat{b}, \lambda) \right| \leq 1 \quad (24)$$

Using relations (22)-(24) and integrating over λ , one can prove (19), in a fashion similar to the proof of CHSH inequality. Based on the relations (21-a) and (21-b), the function $E_{eff}^{(12)}(\hat{a}, \hat{b})$ in (18) has the following relation with the hidden variables level:

$$E_{eff}^{(12)}(\hat{a}, \hat{b}) = \left(\frac{1}{1 - P_0^{(1)}(\hat{a})}\right) \left(\frac{1}{1 - P_0^{(2)}(\hat{b})}\right) \int_{\Lambda} \varepsilon^{(1)}(\hat{a}, \lambda) \varepsilon^{(2)}(\hat{b}, \lambda) \rho(\lambda) d\lambda \quad (25)$$

Solution III. Unlike the first and second solutions, here, we do not make any assumption about the nature of the non-detection probabilities. Instead, we make a conjecture that one can replace (13) by

$$E_{eff}^{(12)}(\hat{a}, \hat{b}) = \int_{\Lambda} \varepsilon_{eff}^{(1)}(\hat{a}, \lambda) \varepsilon_{eff}^{(2)}(\hat{b}, \lambda) \rho(\lambda) d\lambda \quad (26)$$

where $\varepsilon_{eff}^{(1)}(\hat{a}, \lambda)$ and $\varepsilon_{eff}^{(2)}(\hat{b}, \lambda)$ are defined as before and $E_{eff}^{(12)}(\hat{a}, \hat{b})$ is defined in as (18).

One can prove the inequality (19) by using (22)-(24) and (26). The relations (13) and (26) are identical in the ideal limit, but they have different predictions for the real experiments. The physical content of the relation (26) is that *one can always reproduce experimental results using the predictions of a SLHV theory*, whereas relations like (13) indicate that in real experiments one cannot reproduce the predictions of quantum mechanics without making extra assumptions.

Our three solutions for reproducing the inequality (19) involve compatible assumptions. The conjunction of the first two solutions means that the probability of non-detection for a given particle should be merely a function of instrumental efficiencies. Then, the relations (25) and (26) are obtained by dividing both sides of (13) by a detection constant. In such a case, it is assumed that non-detection is only an instrumental problem which is present but does not depend on what a microphysical theory is aimed to describe.

3 The Predictions of Quantum Mechanics

What are the predictions of quantum mechanics for the inequality (19)? In a real double-channel experiment, the respective quantum mechanical joint probability for detecting two photons is nearly equal to [15]:

$$P_{rq, QM}^{(12)}(\hat{a}, \hat{b}) \approx \frac{1}{4} \eta_1 \eta_2 f_{12} \left[1 + r q F \cos 2(\hat{a} - \hat{b}) \right] \quad (27)$$

In this relation, η_k is the efficiency of detecting the k th photon ($k = 1, 2$). The function $f_{12} = f_1 f_2$ shows the probability that both photons reach their detectors, where f_1 denotes the probability for the first photon reaching its corresponding detector and f_2 is the same probability for the second photon. They are indicating the efficiencies of the two corresponding collimators for

photons 1 and 2. The function F is a measure of the correlation of the two emitted photons. In the relation (27), the efficiencies of the analyzers are assumed to be approximately perfect, which is the case in all recent photonic experiments. In an ideal experiment, all of the above efficiencies are equal to one. Here, for simplicity, we assume that $\eta_1 \approx \eta_2 \approx \eta$. Then, using (27), we obtain:

$$\sum_{r,q=\pm 1} P_{rq,QM}^{(12)}(\hat{a},\hat{b}) \approx \eta^2 f_{12} \quad (28)$$

which is independent of polarization directions. Since $(1 - P_{0,QM}^{(1)}) \approx \eta f_1$ and $(1 - P_{0,QM}^{(2)}) \approx \eta f_2$, the relation (28) is also equal to $(1 - P_{0,QM}^{(1)}) (1 - P_{0,QM}^{(2)})$.

Now, using the fact that $E_{QM}^{(12)}(\hat{a},\hat{b}) = \sum_{r,q=\pm 1} rq P_{rq,QM}^{(12)}(\hat{a},\hat{b})$, the quantum correlation function for the polarization directions $\hat{a}$ and $\hat{b}$ is:

$$E_{QM}^{(12)}(\hat{a},\hat{b}) \approx \eta^2 f_{12} F \cos 2(\hat{a} - \hat{b}) \quad (29)$$

and subsequently,

$$E_{QM,eff}^{(12)}(\hat{a},\hat{b}) = \frac{E_{QM}^{(12)}(\hat{a},\hat{b})}{\sum_{r,q=\pm 1} P_{rq,QM}^{(12)}(\hat{a},\hat{b})} \approx F \cos 2(\hat{a} - \hat{b}) \quad (30)$$

If we choose $|\hat{a} - \hat{b}| = |\hat{a}' - \hat{b}| = |\hat{a}' - \hat{b}'| = \varphi$ and $|\hat{a} - \hat{b}'| = 3\varphi$, then (19) yields

$$F |3 \cos \varphi - \cos 3\varphi| \leq 2 \quad (31)$$

For $\varphi = \frac{\pi}{4}$, we have $F\sqrt{2} \leq 1$. In real experiments where the entangled photon pairs are produced through spontaneous parametric down-conversion, F is about 0.95 or more [16, 17]. Since the inequality (31) is independent of the efficiency of detectors and collimators (two main facts responsible for the FS assumption), the predictions of quantum mechanics violate (19) and thus (31), without using the FS assumption. This may be the reason why in spite of the low efficiencies in Bell's photonic experiments, the value of U_{eff} in (19) agrees so well with predictions of the standard quantum mechanics and why this value is nearly the same in different experiments with different efficiency factors.

Appendix

Here, we want to elucidate the meaning of FS assumption more clearly. The CHSH inequality can be expressed as

$$|U| \leq 2 \quad (\text{A-1})$$

where U is a linear combination of some empirical correlation functions along different directions defined in (16). The main issue of FS assumption is that one can use the inequality (19) $|U_{eff}| \leq 2$ instead of (A-1) where U_{eff} is defined in (20).

But, how is it possible to obtain (19) from (A-1) and what is the role of FS assumption in deriving (19)? To answer these two questions, we first remember that an effective correlation function measured in the photonic experiments can be defined as (18). Now, it is obvious that for every $\hat{k} = \hat{a}$ or $\hat{a}'$ and $\hat{l} = \hat{b}$ or $\hat{b}'$, we should have $|E_{eff}^{(12)}(\hat{k}, \hat{l})| \geq |E^{(12)}(\hat{k}, \hat{l})|$, because $\sum_{r,q=\pm 1} P_{rq}^{(12)}(\hat{k}, \hat{l}) \leq 1$. We reformulate $E^{(12)}(\hat{k}, \hat{l})$ as $E^{(12)}(\hat{k}, \hat{l}) = E_{eff}^{(12)}(\hat{k}, \hat{l}) - \epsilon_{kl}$,

where $\epsilon_{kl} := \frac{E^{(12)}(\hat{k}, \hat{l}) \left(1 - \sum_{r,q=\pm 1} P_{rq}^{(12)}(\hat{k}, \hat{l}) \right)}{\sum_{r,q=\pm 1} P_{rq}^{(12)}(\hat{k}, \hat{l})}$. Consequently, one can begin with the CHSH inequality (A-1) and reach the following one

$$-2 + \epsilon \leq U_{eff} \leq 2 + \epsilon \quad (\text{A-2})$$

where $\epsilon = \epsilon_{ab} - \epsilon_{a'b} + \epsilon_{ab'} + \epsilon_{a'b'}$. Since, there is no way to measure $E^{(12)}(\hat{k}, \hat{l})$ in real photonic experiments, the empirical value of ϵ cannot be determined. Considering the predictions of quantum mechanics (see section 3), however, one can show that $\epsilon_{QM} = (1 - \eta^2 f_{12}) U_{eff}$, where η and f_{12} are some efficiencies defined in section 3. Thus, what quantum mechanics predicts is that $|U_{eff, QM}| \leq \frac{2}{\eta^2 f_{12}}$ which is far from violation in actuality.

Yet, there are two situations in which ϵ can be assumed to be zero:

- 1- The experiment is performed under *ideal* conditions, that is $\sum_{r,q=\pm 1} P_{rq}^{(12)}$ reaches one actually.
- 2- The statistics of the experiment can be *fairly* constructed on the basis of the accessible data, that is all the predicted values will remain valid when $\sum_{r,q=\pm 1} P_{rq}^{(12)}$ is renormalized to unity (FS assumption).

There is no way to obtain (19) from (A-1) except for the above conditions. No assumption about the nature of *non*-detection probabilities does help. The renormalization of $\sum_{r,q=\pm 1} P_{rq}^{(12)}$ as well as the validness of the predictions under new conditions are the key points.

Nevertheless, it is logically possible to derive (19) by a *different* approach. One possible way is to begin with an alternative inequality $|U| \leq 2 \sum_{r,q=\pm 1} P_{rq}^{(12)}$ (relation (15)) which is the basis of our first solution. In our two other solutions, we suppose some alternative relations for $E_{eff}^{(12)}(\hat{a}, \hat{b})$ corresponding to the hidden-variable level, to derive (19). So, our three suggested solutions impose more stringent conditions on a SLHV theory for being compatible with the experiments.

References

- [1] J. S. Bell, *Physics* **1**, 195 (1964).
- [2] W. Tittel and G. Weihs, *Q. Inf. Comp.*, **1**, No. 2, 3 (2001).
- [3] M. A. Rowe *et al.*, *Nature* **409**, 791 (2001).
- [4] P. Pearle, *Phys. Rev. D* **2**, 1418 (1970).
- [5] E. Santos, *Phys. Lett. A* **212**, 10 (1996).
- [6] N. Gisin and B. Gisin, *Phys. Lett. A* **260**, 323 (1999).
- [7] J. F. Clauser, M. A. Horne, A. Shimony and R. A. Holt, *Phys. Rev. Lett.* **23**, 880 (1969).
- [8] P. Grangier, *Nature* **409**, 774 (2001).
- [9] E. Santos, *Phys. Lett. A* **200**, 1 (1995).
- [10] J. F. Clauser and M. A. Horne, *Phys. Rev. D* **10**, 526 (1974).
- [11] A. Garg and D. Mermin, *Phys. Rev. D* **35**, 3831 (1987).
- [12] J. Larsson, *Phys. Rev. A* **57**, 3304 (1998).
- [13] A. Shafiee, *J. Mod. Opt.*, **51**, 1073 (2004), quant-ph/0401022.
- [14] A. Shafiee and M. Golshani, *Fortschritte Phys.* **53**, 105 (2005), quant-ph/0406042.
- [15] J. F. Clauser and A. Shimony, *Rep. Prog. Phys.*, **41**, 1881 (1978).
- [16] G. Weihs *et al.*, *Phys. Rev. Lett.* **81**, 5039 (1998).
- [17] W. Tittel *et al.*, *Phys. Rev. A* **59**, 4150 (1999).